# Study of Cronin effect and nuclear modification of strange particles in d-Au and Au-Au collisions at 200 GeV in PHENIX

Dmitri Kotchetkov (*University of California at Riverside*)



for PHENIX Collaboration

### Strangeness at PHENIX

#### **Motivations:**

- ✓ Strange particles as a tool to quantify the effects of medium modification
- ✓ Strangeness observables to look into initial (gluon saturation) or final state (quark recombination, flow)
- ✓ Effects of strangeness on energy loss

#### PHENIX ongoing analyses:

single 
$$K^+, K^-$$

$$K^0_S \to \pi^+ \pi^- \qquad \Lambda \to p \pi^- \qquad \overline{\Lambda} \to p^- \pi^+$$


$$\phi \to K^+ K^- \qquad \phi \to e^+ e^-$$

### Nuclear enhancement and suppression

$$R_{CP} = \frac{\text{Yield(central)} / < N_{coll}(\text{central}) >}{\text{Yield(peripheral)} / < N_{coll}(\text{peripheral}) >}$$

3 0-20%/60-88.5% d+Au π++π-

Parallel Session talk
"π/K/p production and
Cronin effect from p-p,
d-Au and Au-Au
collisions at 200 GeV"
by Felix Matathias



PHENIX d+Au PRELIMINARY

### Mesons vs. baryons or heavier vs. lighter?

#### In central Au-Au collisions:

- ✓ No suppression of protons at  $P_t > 2.0$  GeV
- ✓ Suppression of  $\pi^0$  up to measurement limits (~10 GeV)


#### In central d-Au collisions:

✓ Nuclear enhancement (Cronin) is larger for protons

### How strangeness affects nuclear modification?


- ✓ Effect of strange quarks on R<sub>cp</sub>
- ✓ Strange baryons and antibaryons vs. strange mesons (number of quarks)
- ✓ Mass dependence of  $R_{cp}$  among strange particles

#### **Detectors**



### Hadron's time of flight

In Time of flight Counters (TOF): In Electromagnetic Calorimeter (EMC):



time of flight resolution:

**TOF:** 115 ps


EMC: 700 ps (average) function of energy of a cluster

#### A reconstruction

- √ high asymmetry of decay
- ✓ mean P of  $\pi$  from  $\Lambda$  decay equals 0.3 GeV
- ✓ detect protons in high resolution TOF (up to 3 GeV)
- ✓ reconstruct protons into pairs with any hadron detected either in TOF or EMC
- event mixing technique to build a combinatorial background

### $p\pi$ invariant mass from d-Au collisions

counts/2.5(MeV/ $c^2$ )



invariant mass (GeV/c²)

From 63 x 10<sup>6</sup> minimum bias d-Au collisions:

Λ:

Counts = 24395 + /-373(stat)


Λ–bar:

Counts = 9744 + /-229(stat)

$$\frac{\Lambda:}{S/B} = 1/5$$
  $\frac{S}{\sqrt{S+B}} = 65$ 


$$\Lambda$$
-bar:  $\frac{S}{S/B} = 1/4$   $\frac{S}{\sqrt{S+B}} = 43$ 

counts/2.5(MeV/c<sup>2</sup>)



invariant mass (GeV/c<sup>2</sup>)

### pπ invariant mass from Au-Au collisions



$$\frac{\Lambda:}{S/B} = 1/33 \qquad \frac{S}{\sqrt{S+B}} = 43$$

$$\Lambda$$
-bar: S/B = 1/33

$$\frac{\mathsf{S}}{\sqrt{\mathsf{S}+\mathsf{B}}}=38$$

From 20 x 10<sup>6</sup> minimum bias Au-Au collisions:


Λ:

Counts = 62786 + /-1580(stat)


Λ–bar:

Counts = 48377 + /-1358(stat)

counts /5(MeV/c<sup>2</sup>)




### Detector acceptance normalization



- ✓ Single particle generator ( $K_S^0$ ,  $\Lambda$ , e t.c.)
- ✓ Simulation of PHENIX detector response
- ✓ Extract particle yields as for real data

### $\Lambda$ and $\Lambda$ -bar $P_t$ spectra in d-Au Minimum bias collisions at 200 GeV




Poster Strangeness 5 Arkadij Taranenko


Only statistical errors are shown

### • reconstruction


- $\checkmark \phi \rightarrow K+K- channel$
- √ identify kaons either in TOF or EMC
- event mixing technique to build a combinatorial background

### K<sup>+</sup>K<sup>-</sup> invariant mass from Au-Au collisions





counts/1(MeV/c<sup>2</sup>)  $\frac{\chi^2/\text{ ndf}}{54.83/37}$ 



From 19 x 10<sup>6</sup> minimum bias Au-Au collisions:


 $\phi$ : Counts = 5560+/-240(stat)

S/B = 1/8.5

Posters:
Strangeness 14
by Charles Maguire

Flow 7 by Debsankar Mukhopadhyay

### 



$$\phi \rightarrow K^+K^-$$


Minimum bias events  $dN/dy=1.34\pm0.09(stat)\pm0.20(syst)$   $T=366\pm11(stat)\pm18(syst)$  MeV

0-10% on correct scale, others offset by factors of 10

Parallel Session talk "Light vector mesons (φ) in d-Au collisions in PHENIX" by Richard Seto


### Cronin effect in d-Au collisions

# R<sub>cp</sub> of identified hadrons (0-20% d-Au central collisions) at 200 GeV




Only statistical errors shown for  $\Lambda$ 

# R<sub>cp</sub> of identified hadrons (20-40% d-Au central collisions) at 200 GeV



 $\Lambda$ 's  $R_{cp}$  modification is very similar to one of the proton


# R<sub>cp</sub> of identified hadrons (40-60% d-Au central collisions) at 200 GeV




Mass of  $\Lambda$  is close to one of a proton

### Nuclear modification in Au-Au collisions

# R<sub>cp</sub> of identified hadrons (0-10% Au-Au central collisions) at 200 GeV



## $R_{cp}$ of $\phi$ (0-10% Au-Au central collisions) at 200 GeV



Mass of  $\phi$  is close to one of a proton

### **Summary**


Are differences in R<sub>cp</sub> attributable to mass or quark number?


- ✓ There is no evidence for mass dependence of  $R_{cp}$
- ✓ Strangeness seems to have no effect on R<sub>cp</sub>
- ✓ There is a difference in  $R_{cp}$  for mesons and baryons (see STAR results of  $\Lambda$ 's  $R_{cp}$  in Au-Au)

### Outlook

- ✓  $R_{cp}$  results from  $K^0_S$  and from  $\Lambda$  (Au-Au)
- ✓ Analysis of multi-strange baryons ( $\Xi^0$ ,  $\Xi^+$ ,  $\Xi^-$ ,  $\Omega^-$  and others)

### Extracted K<sup>0</sup><sub>S</sub> signal





From  $48.85 \times 10^6$  minimum bias p-p collisions: Counts = 16630 + /-605(stat)  $62.20 \times 10^6$  minimum bias d-Au collisions: Counts = 116397 + /-2627(stat)